

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

(12) UK Patent Application (19) GB (11) 2 315 753 (13) A

(43) Date of A Publication 11.02.1998

(21) Application No 9714412.5

(22) Date of Filing 08.07.1997

(30) Priority Data

(31) 96305497	(32) 26.07.1996	(33) EP
96305499	26.07.1996	
96308362	19.11.1996	
97301719	14.03.1997	
97301733	14.03.1997	

(71) Applicant(s)

Unilever Plc

(Incorporated in the United Kingdom)

Unilever House, Blackfriars, LONDON, EC4P 4BQ,
United Kingdom

(72) Inventor(s)

Louise Jane Byass
Donald Frank Darling
Charlotte Juliette Doucet
Richard Anthony Fenn
Peter John Lillford
Andrew John McArthur

(51) INT CL⁶

C07K 14/415, A23G 9/00, A23L 3/36

(52) UK CL (Edition P)

C3H HFX HF1
A2B BMF2

(56) Documents Cited

WO 92/22581 A1 US 4565643 A
Biotechnology Advances, Vol. 13, No. 3, 1995, pp
375-402 Plant Physiol., Vol. 104, 1994, pp 971-980
Cryobiology, Vol. 30, 1993, pp 322-328 Advances in
Low Temperature Biology, Vol. 2, 1993, pp 131-182

(58) Field of Search

UK CL (Edition O) C3H HFX HF1
INT CL⁶ A23G 9/00 9/02, A23L 3/36, C07K 14/415
ONLINE: WPI, DIALOG/BIOTECH

(72) cont

David Needham
Christopher Michael Sidebottom
Keith Smallwood
Margaret Felicia Smallwood

(74) continued overleaf

(54) Plant antifreeze proteins for use in frozen confectionary

(57) A frozen confectionery product, e.g. ice-cream comprises one or more antifreeze proteins (AFP's) derived from plants wherein the AFP produces an ice crystal size of less than 15µm after quick freezing to -40°C or less, followed by storage for 1 hour at -6°C. The AFP's can be extracted inter alia from cold-acclimatised, carrots, grass or winter rye.

GB 2 315 753 A

(74) Agent and/or Address for Service

Susan Edith Kirsch

**Unilever Plc, Patent Division, Colworth House,
Sharnbrook, BEDFORD, MK44 1LQ, United Kingdom**

Frozen Food productTechnical Field of the Invention

- 5 The invention relates to frozen food products containing AFPs.

Background to the Invention

- 10 Antifreeze proteins have been described in the literature, see for example Marilyn Griffith and K. Vanya Ewart in Biotechnology Advances, Vol 13, No 3, pp 375-402, 1995. Antifreeze proteins generally possess one or more of the following properties: thermal hysteresis, inhibition of ice
15 recrystallisation, control of ice crystal shape and interaction with ice nucleators.

- Thermal hysteresis is the best known property of AFPs and the property is normally used to test for the presence of
20 AFPs. Thermal hysteresis results from a lowering of the apparent freezing temperature of a solution containing a thermal hysteresis active AFP without affecting the melting temperature. The identification of sources of AFP by thermal hysteresis tests is widely described in the
25 literature, see for example John G. Duman in Cryobiology 30, 322-328 (1993).

- Inhibition of ice recrystallisation is another property of AFPs. This activity is also referred to as ice crystal
30 growth suppression. This property can be tested by comparing at a certain point in time the ice crystal size of crystals in the presence of AFP and in the absence of AFP. The application of this method in the testing of fish AFPs is described in US patent 5,118,792 (DNA Plant
35 Technology Corporation)

A third property of AFPs is their ability to influence the shape of ice crystals. This property stems from the selective binding of AFPs to certain faces of the ice crystal and therewith limiting crystal growth in certain
5 directions. The presence of ice crystals having an hexagonal bipyramid shape is then considered indicative of the presence of AFP. This method is for example described for testing the activity of extracellular winter rye AFPs in WO 92/22581 (University of Waterloo).

10

A fourth property of AFPs is their ability to inhibit the activity of ice nucleating substances. This interaction between an AFP and an ice nucleator may for example result in increased thermal hysteresis. This property is for
15 example tested in WO 96/40973 (University of Notre dame du Lac)

AFPs have been suggested for improving the freezing tolerance of products. Many applications have been
20 suggested in this context.

For example AFPs have been suggested for enhancing the cryopreservation of biological materials (WO 91/12718, Agouron Pharmaceuticals, WO 91/10361, The Regents of the
25 University of California). Also AFPs have been suggested to prevent leakage from liposomes e.g. in cosmetic or pharmaceuticals (see WO 96/20695). A further possible application is to increase the freezing tolerance of plants by including therein (or transgenetically producing
30 therein) an AFP (See J. Cell. Biochem. Suppl. vol. 14e, 1990, page 303 XP002030248, Lee et al, abstract R228). Also fish AFPs have been suggested for use in food products for example in frozen yoghurt or ice cream (US 5,620,732 Pillsbury and WO 96/11586, HSC Research and development
35 limited partnership).

Up till now, however the use of AFPs has not been applied

on a commercial scale. Applicants are of the opinion that one of the reasons for the lack of commercial implementation is that although many AFPs have been described, in practice the implementation in actual commercial products encounters serious problems.

Applicants have found that one of the key reasons for these problems is that out of the great number of AFPs that have been described in the literature only a limited set of AFPs can suitably be applied for each application; also applicants have found that this selection of suitable AFPs is dependent on the desired application and/ or product attributes to be achieved.

A particular desirable source of AFPs is plant material. Plant materials can fairly easily be obtained in relatively large quantities and relatively simple isolation procedures can be used for obtaining an AFP containing concentrate. Furthermore the use of AFPs from plant material is believed to be favoured by consumers who tend to prefer natural vegetable sources to e.g. fish AFPs.

Marilyn Griffith and K. Vanya Ewart in Biotechnology Advances, Vol 13, No 3, pp 375-402, 1995 have given a list of 27 higher plant species in which antifreeze activity is found. This article also suggests a wide range of possible applications for AFPs.

Applicants have now found that if a specific application for the use of plant AFPs is selected, this creates the need for a specific test to select the limited set of AFPs which can advantageously be applied in this application.

The object of the present invention is therefore to provide those plant AFPs which can advantageously be used in frozen confectionery products.

- 5 Surprisingly, applicants have found that despite the fact that a great number of plants contain AFPs, only a limited set of plants contains AFPs which are capable of providing a good texture to frozen confectionery products. Surprisingly it has been found that a relatively simple
10 test method can be used to select the suitable AFPs.

Accordingly in a first aspect the invention relates to frozen confectionery products comprising one or more AFPs derived from plants, wherein the AFPs in water have an ice
15 crystal size after quick freezing to -40°C followed by storage for 1 hour at -6°C (measured as described below) of less than $15\text{ }\mu\text{m}$.

20 Background to the invention

A number of literature places have suggested that AFPs may potentially be used for favourably influencing the textural properties of frozen confectionery products such as ice
25 cream. However most of these documents do not provide a teaching how these favourable properties can actually be achieved in practice.

A further set of documents describes the use of fish AFPs.
30

WO 96/11586 teaches the application of fish antifreeze polypeptides in frozen fermented food products. This document does not teach the use of specific AFPs derived from plants in these products.

35

WO 96/39878 describes the application of AFP in ice-cream. Suitable AFPs for this application may be derived from

blood and muscle tissue of antartic fish, arttic fish, worms and insects. Again no teaching is provided that plant AFP can be used.

- 5 US 5,118,792 describes in example 3B the inhibition of recrystallisation by a purified A-Saf5 fusion protein in a popsicle mixture. Again this document does not teach the use of AFPs derived from plants.
- 10 WO 92/22581 describes a plurality of polypeptides derived from the extracellular spaces of winter rye. Several possible applications of these polypeptides are described in general, among these is ice-cream. However no teaching is provided which of the polypeptides should be selected to
- 15 obtain a good quality ice-cream. Applicants have found that only specific proteins from winter rye are suitable for use in ice-cream (see examples).
- Applicants have found that a great number of plants which
- 20 up till now have not been listed as such, contain a significant level of AFPs. On the other hand it has been found that not all AFP containing plants provide the right type of AFP for favourably influencing the texture of frozen confectionery. Applicants now aim at providing a
- 25 particular novel selection of plant sources which surprisingly provide on the one hand good AFP properties and on the other hand are capable of favourably influencing the textural properties of ice cream.
- 30 In particular it has been found that suitable AFPs can be selected by taking a composition comprising AFP in an aqueous composition, quick freezing this composition to -40 or less followed by storage for 1 hour at -6 °C. Suitable AFPs result in an ice crystal size after storage for 1 hour
- 35 at these conditions of less than 15 μm , more preferably 5-14 μm , most preferred 8 to 12 μm . The temperature of quick freezing is advantageously -40 to -100 °C, preferably -80 °C.

A detailed description of a suitable test to determine this characteristic is given in example I.

Generally the test can be applied to any suitable
5 composition comprising AFP and water. Generally the level
of AFP in such a test composition is not very critical and
can for example be from 0.0001 to 0.5 wt%, more preferred
0.0005 to 0.1 wt%, most preferred 0.001 to 0.05 wt%, for
example 0.01 wt%. The water level in the aqueous
10 composition is advantageously 30 wt% or more, for example
50 wt% to 99.9999 wt%

Any suitable composition comprising AFP and water can be
used to carry out the test. If desired additives may be
15 present, e.g. sucrose or buffering agents. Generally,
however, it will not be necessary to obtain the AFP in
purified form. For practical applications normally it would
suffice to prepare a liquid extract or juice of plant
material, wherein this extract or juice can then be tested.
20 A suitable method to prepare suitable liquid compositions
is given in example II.

Normally plants to be tested for suitable AFPs have been
subjected to cold. For example plants can be tested which
25 grow in cold climates, for example antartic plants.
Alternatively plants can be harvested during the winter
period, preferably December to March, more preferred
January to February, most preferred in January (northern
hemisphere) or June to September, more preferred July to
30 August, most preferred July (southern hemisphere).

Applicants have subjected a great number of plants to the
above described test. The results are given in examples III
and IV.
35

Preferred sources of AFPs are derived from *Polystichum*
mohriodes, *Ranunculus biternatus*, *Nothofagus antartica*,

- Cerastium fontanum, Colobanthus quitensis, Rumex acetosella, Salix fragilis, Calluna vulgaris, Aceana magellanica, Pisum sativum, Acer sacchararoides, Oxalis, Geranium, Daucus carota (carrot), Vinca minor (periwinkle),
- 5 Vinca major, Polemonium, Buddleia, Forsythia, Sambucus nigra, Juncus squarrosus, Carex aquatilis, Agrostis tenuis, Deschampsia antartica, Festuca contracta, Festuca rubra, Parodiochloa flabellata, Phleum alpinum, Poa annua (speargrass), Poa pratensis (Kentucky blue grass),
- 10 Rostkovia magellanica, Bambosoideae, Chorisodontium aciphyllum, Drepanocladus uncinatus, Isothenicium myosuriodes, Polytrichum alpestre, Alectoria nigricans, Caloplaca regalis, Himantormia lugubris, Hypogymnia physodes, Parmelia subrudecta, Ramalina farinaceae,
- 15 Stereocaulon glabrum, Umbilicaria antartica, Usnea subfloridana, Poa trivialis, Lolium perenne, Holcus lanatus, Bromus sterilis and Festuca contracta.

Applicants are of the opinion that based on the guidelines

20 given in the description of the invention the skilled person will be well able to select further AFPs which can suitably be derived from plants. The use of these AFPs is also embraced within the scope of the present invention.

25 In one embodiment particularly useful for application in food products are those plant AFP, which satisfy the above defined crystal size test and which are derived from non- or low-toxic plants. In this context preferred are AFP derived from carrots, grasses, bamboo etc.

30

Another preferred selection from the above sources is the AFPs derived from the Lichen family, in particular Alectoria nigricans, Caloplaca regalis, Himantormia lugubris, Hypogymnia physodes, Parmelia subrudecta,

35 Ramalina farinaceae, Stereocaulon glabrum, Umbilicaria antartica, Usnea subfloridana. These AFPs show particularly good properties in frozen confectionery products.

Other AFPs which show particularly good activity are derived from *Juncus squarrosus* or *Geranium*.

- For some applications those AFPs are selected which
- 5 maintain their ability to limit ice crystal growth (as evidenced by the above test) even after heat treatment above a temperature of 60°C, most preferably from 80 to 105 °C for a period of at least 30 seconds, more preferred more than 1 minute, 10 minutes or even more than 1 hour.
- 10 Suitable plant sources which are heat stable are for example *Acer saccharoides*, *Bamboo*, *Buddleia*, *Isothecium myosuroides*, *Ramalina farinaceae*, *Usnea subfloridana*, *Forsythia*, *Oxalis*, *Poa trivialis*, *Lolium perenne*, *Holcus lanatus*, *Bromus sterilis*, *Parodiochloa flabellata*,
- 15 *Deschampsia antarctica*, *Carex aquatilis*, *Colobanthus quintensis* and *Agrostis tenuis*, *Festuca contracta*, *Poa annua*.

- In addition to the plants mentioned above applicants have
- 20 tested the AFPs from *Secale cereale* (winter rye). Several winter rye AFPs have been described in WO 92/22581. Surprisingly applicants have found that the 32 kDa protein derived from winter rye satisfies the above described crystal size test, while other proteins from winter rye do
- 25 not satisfy the test (see example IX.) For the purpose of the invention, therefore if AFPs derived from winter rye are used, preferably the 32 kDa protein is applied.

- Another preferred embodiment of the invention relates to
- 30 the use in frozen confectionery products of plant AFPs which satisfy the above test and which are not derived from winter rye.

Detailed description of the invention

35

Frozen products in accordance to the present invention comprise at least one AFP, which can be derived from plant

sources.

The AFPs can be obtained from the plant sources by any suitable process, for example the isolation processes as described in the above mentioned documents. Alternatively the AFPs can be extracted from the plants, for example by preparing a plant extract from (parts) of the plant followed by an optional concentrating step. Examples of suitable methods for obtaining said extracts are given in the examples.

Also microorganisms or plants may be genetically modified to express AFPs and the AFPs may then be used in accordance to the present invention.

Genetic manipulation techniques may be used to produce AFPs having at least 80%, more preferred more than 95%, most preferred 100% homology to the AFPs directly obtained from plant sources which naturally contain the AFPs. For the purpose of the invention these AFPs possessing this high level of homology are also embraced within the term "AFP derived from plants". For the purpose of the invention the term "AFP derived from plants" preferably does not include AFPs which naturally occur in non-plant sources e.g. fish and which by transgenetic routes are produced by plants.

The genetic manipulation techniques may be used as follows: An appropriate host cell or organism would be transformed by a gene construct that contains the coding region for the desired polypeptide.

The nucleotide sequence coding for the polypeptide can be inserted into a suitable expression vector. Said vector encoding the necessary elements for transcription and translation, and in such a manner that they will be expressed under appropriate conditions (e.g. in proper orientation and correct reading frame and with appropriate

targeting and expression sequences).

The methods required to construct these expression vectors are well known to those skilled in the art.

5

A number of expression systems may be utilised to express the polypeptide coding sequence. These include, but are not limited to, bacteria, yeast, insect cell systems, plant cell culture systems and plants all transformed with the appropriate expression vectors.

10

AFPs obtainable from the above mentioned sources can be used in any suitable frozen confectionery product. For the purpose of the invention the term frozen confectionery product includes milk containing frozen confections such as ice-cream, frozen yoghurt, sherbet, sorbet, ice milk and frozen custard, water-ices, granitas and frozen fruit purees. For some purposes the use in fermented frozen food products is less preferred.

15

Preferably the level of AFPs in the frozen confectionery product is from 0.0001 to 0.5 wt% based on the final product, more preferred 0.0005 to 0.3 wt%, most preferred 0.001 to 0.2 wt%.

20

Preferably the level of solids in the frozen confection (e.g. sugar, fat, flavouring etc) is more than 2 wt%, more preferred from 4 to 70wt%.

30 If desired the frozen confectionery products of the invention may be aerated, for example to an overrun of from 50 to 500%.

35 The method of preparing the frozen confectionery product of the invention can be selected from any suitable method for the preparation. The AFPs can generally be added at various stages of the preparation, for example it can be added in

the first pre-mix of ingredients or can later be added during a later stage of the preparation process. For some applications it is sometimes preferred to add the AFPs at a relatively late stage of the production process, for
5 example after (partial) pre-freezing of the product.

The freezing process for frozen confectionery products can be selected from any suitable freezing process and may optionally comprise an aeration step for example to an
10 overrun of 50 to 300 %. For some purposes it is advantageous that the freezing process involves a cold hardening step, for example at a temperature of -30 Fahrenheit or lower.

15 For some applications it may be advantageous to include a mixture of two or more different AFPs into the frozen confectionery product. One reason for this can for example be that the plant source for the AFP's to be used, contains more than one AFP and it is more convenient to add these,
20 for example because they are both present in the plant extract to be used. Alternatively it may sometimes be desirable to add more than one AFP from different sources.

The invention will now be illustrated by means of the
25 following examples

Example I

Test for determining ice crystal particle size after quick cooling followed by storage at -6 °C for 1 hour.

5

The preferred method is as follows:

Ia: Anti-freeze activity was measured using a modified "splat assay" (Knight et al, 1988). 2.5 µl of the solution under investigation in 30% (w/w) sucrose was transferred onto a clean, appropriately labelled, 16 mm circular coverslip. A second coverslip was placed on top of the drop of solution and the sandwich pressed together between finger and thumb. The sandwich was dropped into a bath of hexane held at -80°C in a box of dry ice. When all sandwiches had been prepared, sandwiches were transferred from the -80°C hexane bath to the viewing chamber containing hexane held at -6°C using forceps pre-cooled in the dry ice. Upon transfer to -6°C, sandwiches could be seen to change from a transparent to an opaque appearance. Images were recorded by video camera and grabbed into an image analysis system (LUCIA, Nikon) using a 20x objective. Images of each splat were recorded at time = 0 and again after 60 minutes.

25

Alternatively (less preferred) the properties can be measured as follows:

Ib: A sample of an AFP containing product containing water is adjusted to a sucrose level of 30 wt% (If the starting level of the sample is more than 30% this was done by dilution, if the starting level was lower sucrose was added to the 30% level).

35 A 3 µL drop of the sample is placed on a 22 mm coverslip. A 16 mm diameter cover-slip is then placed on top and a 200 g weight is placed on the sample to ensure a uniform slide

thickness. The edges of the coverslip are sealed with clear nail varnish.

The slide is placed on a Linkham THM 600 temperature
5 controlled microscope stage. The stage is cooled rapidly
(50 °C per minute) to -40°C to produce a large population
of small crystals. The stage temperature is then raised
rapidly (50°C per minute) to -6 °C and held at this
temperature.

10

The ice-phase is observed at -6 °C using a Leica Aristoplan
microscope. Polarised light conditions in conjunction with
a lambda plate were used to enhance the contrast of the
ice-crystals. The state of the ice phase (size of ice-
15 crystals) is recorded by 35 mm photomicrography at T=0 and
T=1 hour. Whereby an average particle size (visual
determination, number average) of below 15 µm indicates a
suitable AFP for use in frozen confectionery products.

Example II

Methods to obtain compositions containing AFP and water.

- 5 A. Fresh tissue of plant parts, for example roots, stems, buds or leaves can ground with a pestle and mortar (cooled to 4°C) in an equal volume buffer A (for example 10mM EDTA, 20 mM Ascorbic acid, buffered with Tris to pH 7.4) held on ice. The homogenates are filtered through one or
10 more layers of muslin and kept on ice prior to further use.

This method can generally be applied to most plants and provide fresh plant juice containing the AFP. Generally the entire plant can be used for this purpose although for
15 practical reasons only parts may sometimes be used (for example leaves from woody trees, roots from root vegetables and stems from grassy plants).

- B. Method to extract AFPs from sources which are capable of
20 withstanding heat such as grasses. This method is exemplified by using mixed grass. However it will be apparent that this method may equally be applied to other heat stable sorts.

- 25 Mixed grass tissue (*Poa Trivialis*, *Lolium Perenne*, *Holcus Lanatus*, *Bromus Sterilis*) was cut in January (mean temperature in that month was 3.5 °C ensuring the appropriate cold acclimatization of the plants). The grass tissue was rapidly transported into the laboratory for
30 further handling and washed thoroughly with water to remove dirt.

- 500 g of grass clippings was placed in a 650 Watt microwave oven and heated at full power for 5 minutes, whereby the
35 temperature was raised to 85 to 100°C. The grass clippings were then cooled to ambient temperature.

After the heating step the AFP rich juice was separated from the clippings by filtering. The mass was stirred continuously for 5 minutes in the presence of an equal volume of water and then squeezed through 3 layers of 5 muslin.

Example III

Screening of various plants. Non antarctic plants were harvested in January (mid winter). The antarctic plants
5 were harvested mid summer (February-March).

Unless indicated otherwise roots were used to prepare an AFP containing juice according to the method as described in example IIa.

10

The samples were subjected to the test of Example Ia. Suitable AFPs for application in frozen confectionery products are indicated with a positive sign (+).

- 15 Additionally the thermal hysteresis properties of the juices was measured as follows: The thermal hysteresis of a sample of a product containing AFPs was determined by placing the melted product on a microslide (Camlab Cambridge, path length 0.1 mm). The ends of the microslide
20 are sealed with petroleum jelly. Ice is introduced into the sample using an aerosol freezing spray. The slide was then immersed in ethanol temperature regulated bath at -0.1°C . After 5 minute equilibration the sample is checked. if the ice melts completely the temperature of the bath was
25 lowered in 0.1°C steps followed by equilibration. These steps are repeated until a temperature was reached where a small amount of ice crystals exist in the sample. After equilibration at that temperature, the bath temperature was decreased in steps of 0.01°C per minute. The freezing
30 point of the sample is recorded as the temperature at which the ice propagation begins from the equilibrated crystals. The melting temperature of the sample is then determined by raising the temperature starting at the freezing point in steps of 0.01°C per minute until all ice-crystals melt.
35 This temperature is the melting temperature of the sample. The thermal hysteresis of the sample is the difference between the melting temperature and the freezing

temperature. AFPs with a significant degree of thermal hysteresis are indicated with a positive sign (+). The following results are obtained:

5	Plant name	Thermal Hysteresis	Recrystallisation inhibition (Ex I)
	<i>Equisitum hymenale</i>	+	-
	<i>Picea glauca</i>	+	-
	<i>Dicentra cucularia</i>	+	-
10	<i>Viola</i> sp.	+	-
	<i>Brassica oleracea</i>	+	-
	<i>Brassica rapa</i>	+	-
	<i>Brassica napus</i>	+	-
	<i>Daucus carota</i>	+	+
15	<i>Vinca minor</i>	+	+
	<i>Solanum tuberosum</i>	+	-
	<i>Poa annua</i>	+	-
	<i>Poa pratensis</i>	+	-
	<i>Secale cereale</i>	+	*)
	<i>Petroselinum crispum</i>	+	-

Salvia officinales	+	-
--------------------	---	---

*) Note: active but only the 29-32 kDa protein

- 5 These results clearly show that although a great number of plants show thermal hysteresis properties, only a small proportion of these satisfy the recrystallisation test.

Example IV

Screening of various plants. Non antarctic plants were harvested in January (mid winter). The antarctic plants
5 were harvested mid summer (February-March).

Unless indicated otherwise roots were used to prepare an AFP containing juice according to the method as decribed in
example IIa.

10

The samples were subjected to the test of Example Ia. Suitable AFPs for application in frozen confectionery products are indicated with a positive sign (+).

15

20

25

PLANT NAME	RI test
Water horsetail	-
Asplenium scolopendrium	-
Polystichum mohriodes	+
Pinus sylvestris	-
Picea glauca	-
Cupress sp	-
Laurus	-
Ranunculus bitermatus	+
Dicentra	-
Platanus orientalis	-
Urtica	-
Pterocarya fraxinifolia	-
Nothofagus antarctica	+
Nothofagus oblique	-

5	Betula pendula	-
	Beta vulgaris	-
	Cerastium fontanum	+
	Colobanthus quitensis (pearlwort)	+
	Rumex acetosella	+
10	Paeonia	-
	Hypericum	-
	Alcea	-
	Viola	-
	Populus alba	-
15	Salix alba 'Britzensis'	-
	Salix daphnoides	-
	Salix fragilis	+
	Sorbus aria	-
	Prassica napus	-
20	Erysimum	-
	Calluna vulgaris	+
	Primula	-
	Hydrangea	
	Sedum	-
25	Acaena magellanica	+
	Crategus monogyna	-
	Cotoneaster spp x2	-
	Fragaria x ananassa	-

	Alchemilla	-
	Cytisus	-
	Pisum sativum	+
	Vicia faba	-
5	Medicago sativa	-
	Daphne	-
	Eucalyptus	-
	Aucuba	-
	Ilex	-
10	Acer saccharoides	+
	Rhus	-
	Oxalis	+
	Geranium	+
	Hedera	-
15	Daucus carota	+
	Pastinaca sativa	-
	Vinca	+
	Polemonium	+
	Rosmarinus	-
20	Buddleia	+
	Forsythia	+
	Fraxinus ornus	-
	Lonicera pileata	-
	Sambucus nigra	+
25	Lactuca sativa	-

	Tragopodon porrifolius	-
	Helianthus tuberosus	-
	Juncus squarrosus	+
	Carex aquatilis	+
5	Agrostis tenuis	+
	Deschampsia antarctica	+
	Festuca contracta	+
	Festuca rubra	+
	Parodiochloa flabellata	+
10	Phleum alpinum	+
	Poa annua	+
	Poa pratensis	+
	Rostkovia magellanica (grass)	+
	Bambosoideae sp	+
15	Muscari armenicum	-
	Allium ampeloprasum cv Alaska	-
	Allium cepa	-
	Chorisodontium aciphyllum	+
	Drepanocladus uncinatus	+
20	Isothecium myosuroides	+
	Neckera complanata	-
	Polystichum alpestre	+
	Polytrichum commune	-
	Polytrichum formosum	-
25	Racomitrium lanuginosum	-

5	Sphagnum capillofolium	-
	Sphagnum palustre	-
	Alectoria nigricans	+
	Caloplaca regalis	+
	Himantormia lugubris	+
10	Hypogymnia physodes	+
	Parmelia subrudecta	+
	Ramalina farinaceae	+
	Stereocaulon glabrum	+
	Umbilicaria antarctica	+
	Usnea subfloridana	+

Example V

A liquid pre-mix for preparing ice-cream was made by mixing:

5		
	Ingredient	% by weight
	Skimmed milk powder	11.390
	Sucrose	3.410
	Maltodextrine (MD40)	4.000
10	Locust bean gum	0.072
	Corn Syrup 63DE	20.705
	Guar Gum	0.048
	Genulacta L100	0.020
	Butter	9.015
15	Avicel RC581	0.240
	Gelatin	0.140
	Monoglyceride (palmitate)	0.450
	Vanillin	0.010
	AFP (of example IIb)	0.100 or none (control)
20	Water	balance

*Note: AFP is added as concentrated AFP solution using some of the added water as a diluent, percentage refers to amount of AFP.

25

This mix can conveniently be pasteurised at 85°C for 15 seconds and stored chilled in a can.

30 The mixes can be used in the preparation of a ice-cream by whipping with a conventional house-hold mixer to an overrun of about 100%, followed by quiescently freezing into a house-hold freezer.

35 The composition according to the invention had a markedly better texture than the control sample.

Similar results can be obtained by using the following

plants sources: Acer saccharoides, Bamboo, Buddleia,
Isothecium myosuroides, Ramalina farinaceae, Usnea
subfloridana, Forsythia, Oxalis, Poa trivialis, Lolium
perenne, Holcus lanatus, Bromus sterilis, Parodiochloa
5 flabellata, Deschampsia antarctica, Carex aquatilis,
Colobanthus quintensis and Agrostis tenuis, Festuca
contracta, Poa annua

Example VI

A liquid premix for the preparation of ice-cream was prepared by mixing:

5	Ingredient	% by weight
	Skimmed milk powder	10.00
	sucrose	13.00
	maltodextrine (MD40)	4.00
10	Locust bean gum	0.14
	butteroil	8.00
	monoglyceride (palmitate)	0.30
	vanillin	0.01
	AFP (of example IIb)	0.01 or none(control)
15	water	balance

*Note: AFP is added as concentrated AFP solution in some of the water, percentage refers to amount of AFP.

20

The ingredients were mixed at ambient temperature followed by pasteurisation for 60 seconds at 89°C. The mix was aseptically filled into packs of 500 ml, sealed and stored at ambient temperatures.

25

The mix can be used for the preparation of ice-cream by whipping it with a conventional house-hold mixer to an overrun of about 70 % followed by freezing under quiescent conditions in a house-hold freezer. After two months

30 storage the composition according to the invention had a markedly better texture than the control sample.

Example VII

This example describes the isolation and sequencing of
5 carrot AFP. Similar methods can be used for other plant
AFPs.

Carrot root tissue of cold acclimatised carrots was
homogenized in three volumes (w/v) buffer (20mM ascorbic
10 acid, 10 mM EDTA, 50 mM Tris/HCL, pH 7.2) in a pre-cooled
pestle and mortar and filtered through one layer of muslin.
The filtrate was centrifuged at 6,000g, ten minutes at 4°C;
the supernatant was collected and centrifuged at 100,000g
for 1 hour at 4°C. The 100,000g supernatant from this step
15 is termed the soluble fraction and the pellet the
microsomal fraction.

The supernatant was applied to a 30 ml fast flow Q
Sephacrose (Pharmacia) column pre-equilibrated in 50 mM
20 Tris/HCL pH 7.4 at a flow rate of 5 ml/min supplied by a
HiLoad pump P-50 controlled by a Gradifrac low pressure
chromatography system (Pharmacia) at 4°C and the eluate
monitored at OD 280 by a UV monitor (Monitor UV1,
Pharmacia) recorded on a chart recorder (REC 102,
25 Pharmacia). 5 ml fractions were collected. The column was
washed with 50mM Tris/HCL pH 7.4 at the same flow rate
until the OD 280 returned to zero. A 150ml gradient of 0-
0.4 M NaCl in Tris/HCL pH 7.4 was then applied followed by
a 2 M NaCl column wash. Eluate fractions were subjected to
30 the splat assay as in example I.

Fractions containing anti-freeze activity were pooled and
concentrated using polyethylene glycol as follows: the
fractions were transferred in 10kDa cut off dialysis tubing
35 (Sigma) which had been washed in tap water, boiled in 50mM
EDTA pH 7.5 for 10 minutes and rinsed in milli Q water. The
dialysis tubing containing the sample to be concentrated

was covered with solid polyethylene glycol compound Mol. Wt. 15,000 - 20,000 (Sigma) and incubated at 4°C for up to 4 hours or until the sample volume inside the dialysis tubing had reduced up to 10 fold.

5

The pooled concentrate from the Q sepharose column was applied either to a phenyl Sepharose column, a SMART superdex 75 gel permeation column or an FPLC superdex 75 gel permeation column.

10

Carrot root anti-freeze proteins were purified by gel permeation chromatography as follows:

20µl aliquots of sample were applied to a SMART superdex 75
15 column (Pharmacia) pre-equilibrated in 50mM Tris/HCl pH7.4 containing 0.15M NaCl (Buffer E) at a flow rate of 40µl/min and components separated by gel permeation at the same flow rate in equilibration buffer. The eluate was monitored at OD 280 and OD 215. 80µl fractions were collected between
20 0.85 and 0.89ml, 40µl fractions between 0.89 and 1.24ml and 100µl fractions between 1.24 and 3.0 ml. The void volume (Vo) of the column was 0.91 ml as determined by the retention volume of a solution of Blue Dextran. The superdex column was calibrated by application of 10µl of a
25 solution containing 5mg/ml BSA (Mr 66kDa, retention (Ve)=1.02 ml), 3mg/ml Carbonic anhydrase (Mr 29 kDa, Ve=1.22 ml), 2mg/ml Cytochrome C (Mr 12.4 kDa, Ve=1.41 ml) and 2mg/ml Aprotinin (Mr 6.5 kDa, Ve=1.59 ml) and a standard curve plotted of Ve/Vo against log Mr. Fractions
30 containing anti-freeze activity were identified by the spat assays in Example I, with an activity peak that showed a retention volume of 1.16 ml and an apparent molecular weight of 40 kDa. These measurement confirmed that the 38 kDa band from cold acclimatised carrots was an anti-freeze
35 peptide.

SDS-PAGE was carried out according to Laemmli (1970) using

- the Biorad mini system. Samples to be analyzed by SDS-PAGE were dissolved in SDS-PAGE sample buffer (Laemmli 1970), heated for 5 minutes at 100°C on a dry heating block (Techne) and centrifuged for 3 minutes at 10,000g at room temperature. Samples (10-50µl) were applied to mini-gels (Biorad, 0.75, 1.0 or 1.5mm thickness, 10, 12, 15% acrylamide or 10-20% gradient acrylamide {pre-poured from Biorad}) and electrophoretically separated. Separated polypeptides were fixed and stained in the gel either with Coomassie blue (0.1% {w/v} Coomassie Brilliant Blue in acetic acid/methanol/miliQ water {5:4:31, by vol}) or silver stained using the Biorad silver stain kit according to the manufacturer's instructions. Gels were dried between two sheets of Gelair collophane in a Biorad gelair dryer according to the manufacturer's instructions. Sigma high and low range molecular weight marker kits were used according to the manufacturer's instructions for determination of apparent M_r on SDS-PAGE.
- 20 The ion exchange chromatography was carried out with cold acclimatized carrot root and non-cold acclimatized carrot root. The resulting gel SDS-PAGE gels showed the presence of a about 38kDa band in the cold acclimatized sample. This band was much less abundant in the non-cold acclimatized root. This (about) 38kDa band was hence attributed to anti-freeze activity.

For protein sequencing, the carrot root protein of about 38kDa was purified as described in the previous example and then to ensure further purification the sample to be sequenced was excised from the SDS PAGE gel and then proteolytically digested *in situ* in the polyacrylamide gel slice.

- 35 Preparations of largely pure of protein about 38 kDa, that still had some minor contaminating proteins, were loaded onto a 12% polyacrylamide gel. Three lanes each with 2 µg

of protein were loaded and electrophoresed in the gel until the dye front reached the bottom of the gel. The gel was then stained in 0.2% coomassie brilliant blue (w/v), 30% methanol (v/v), 1% acetic acid (v/v) for 20 minutes and
5 then destained with 30% methanol until the protein bands could be visualised. The 38 kDa band was identified by comparison with molecular weight markers loaded into adjacent lanes and the band from each lane was excised with a scalpel blade, taking care to exclude contaminating
10 bands.

The gel slices were transferred to a clean eppendorf tube and washed twice with 0.5ml of 50% acetonitrile (v/v), 100mM Tris/Cl, pH 8.5. The washing removed some of the
15 coomassie stained and also partially dehydrated the gel slices. The gel slices were then removed from the tube and subjected to air drying on the laboratory bench until they had shrunk significantly and started to curl up. They were then transferred back to the eppendorf and rehydrated with
20 firstly, 10µl of 100mM Tris/Cl, pH 8.5 containing 1µg of endoproteinase Lys C (Boehringer Mannheim). This is a proteinase that specifically cleaves polypeptide chains on the carboxy terminal side of lysine residues. Further Tris buffer was added to the gel slices until they were fully
25 rehydrated and they were then incubated at 37°C for 16 hours.

After incubation 1µl of trifluoroacetic acid was added to the tube to stop the reaction and then the gel slices were
30 washed twice with 0.3ml of 60% acetonitrile (v/v), 0.1% TFA (v/v) at 30°C for 30 minutes. This was to again partially dehydrate the gel slices causing them to shrink and elute the peptides that had been generated. The supernatant was transferred to another clean eppendorf tube and then dried
35 in a centrifugal

evaporator for 2 hours until the sample was near dryness and resuspended to a volume of 0.1ml with 0.1% TFA.

The peptides were then separated by reversed phase HPLC on
5 a Smart micropurification system (Pharmacia). The peptide
digest was loaded onto a C18 column (2.1 x 100 mm)
equilibrated in 0.1% TFA (Solvent A) at a flow rate of
0.1ml min. The column was then eluted with a gradient of 0
- 70% of Solvent B (90% acetonitrile v/v, 0.085% TFA v/v)
10 over 70 minutes at the same flow rate. The optical density
was monitored at 214 nm and individual peptide peaks were
collected in the fraction collector by manual stepping.
Polypeptides were sequenced by loading onto a model 492
Perkin Elmer protein sequencer using the liquid phase
15 chemistry cycles as recommended by the manufacturer.

Several polypeptide fragments (A-E) were analyzed in the 38
kDa band and had sequences substantially homologous to:

- (A) LEU-PRO-ASN-LEU-PHE-GLY-LYS
- 20 (B) ILE-PRO-GLU-GLU-ILE-SER-ALA-LEU-LYS
- (C) LEU-THR-X-LEU-ASP-LEU-SER-PHE-ASN-LYS
- 25 (D) SER-LEU-ARG-LEU-SER-SER-THR-SER-LEU-SER-GLY-PRO-VAL-
PRO-LEU-PHE-PHE-PRO-GLN-LEU-X-LYS
- (E) X-X-GLU-VAL-ILE-PRO-X-GLN-LEU-SER-THR-LEU-PRO-ASN-LEU-
LYS

30

Cell cultures for producing anti-freeze proteins were made
as follows:

New cell cultures were initiated based on methods described in Gamborg and Wetter 1975, Torres 1989, Dodds and Roberts 1985.

- 5 Cold acclimatised Carrot (Autumn King): the surface of the storage root was sterilized firstly by washing with 10% Teepol detergent, followed by scrubbing under running water then rinsing under running water for 15 minutes. Where practicable (on the basis of size) the root was peeled. The
- 10 root was then aseptically cut into 0.5cm slices, which were placed in 70% v/v ethanol for 10 minutes with shaking followed by 10% v/v Domestos + 2 drops Tween 20 (Sigma) for 25 minutes, also with shaking. Sections were then washed 3 x with sterile distilled water. Cylinders of approximately
- 15 0.5cm diameter were cut through the slices using a sterile scalpel, and the remainder cut into 2-3mm lengths. These tissue discs (explants) were aseptically transferred onto solid MS medium containing 30g/l sucrose, 10mg/l indole acetic acid (IAA), 0.1 mg/l kinetin and 8g/l technical
- 20 agar, which was contained in 60ml Sterilin containers. The explants were incubated in the dark at 20°C.

When necessary, the resulting calli were divided into smaller sections, which were plated onto fresh medium.

- 25 Suspension cultures were then initiated from actively growing callus.

- Additionally carrot cell suspension culture lines (NOR and OX6) were obtained from the Department of Biochemistry and
- 30 Molecular Biology, University of Leeds. 10 ml of these cultures were sub-cultured into 90ml of fresh Murashige and Skoog medium (Sigma) containing 25g/l sucrose and 1mg/l 2,4-D every seven days. Cultures were

incubated in an orbital shaking incubator at 150 rpm at 25°C in the dark.

The NOR culture was cold treated as follows:

- 5
18x5ml 7d old NOR culture was added to 18 x 100ml
Erlenmeyer flasks containing 45ml carrot MS medium. The
cultures were incubated at 25°C as previously described for
4 days, then the incubator temperature reduced to 4°C. Two
10 flasks were removed immediately and the cells and medium
harvested as previously described as $t = 0$. The remaining
flasks were harvested in duplicate at
 $t=8h, 1d, 2d, 4d, 7d, 9d, 11d$, and 14d.
- 15 The cold acclimation treatment was repeated using larger
cultures of both NOR and OX6, which were transferred to 4°C
after 4d and 7d of growth at 25°C. Cultures were harvested
at $t=0$, $t=7d$ and $t=14d$. In addition to harvesting, the PCV
was determined for each culture at each time point.
- 20
The NOR cold acclimated cells were prepared for splat
analysis as in Example I as follows: Quick frozen cells
were ground to a fine powder in liquid nitrogen using a
pestle and mortar. The powdered samples were resuspended in
25 2 x volume of 10mM EDTA + 20mM ascorbic acid, whirlimixed
for 30 seconds then centrifuged at 10.000g for 10 minutes.
10µl aliquots of the supernatants were splatted using the
buffer control as a negative control. RI activity could be
detected in cold acclimated cells and medium but not in the
30 non-cold acclimated samples.

The media samples from NOR suspension were analyzed as
follows. The NOR carrot medium was buffered by addition of
100µl of 1M Tris/HCl pH 7.4. This was then applied to the 1
35 ml Q Sepharose column (Pharmacia) at a flow rate of 1
ml/min and bound molecules eluted with 3 ml aliquots of 50
mM Tris/HCl pH 7.4 containing concentrations of 0.5 M NaCl.

1 ml fractions were collected.

This anion exchange method was also used to fractionate
t=0,2d,4d,7d,11d, cold acclimated and t=7d non-cold
5 acclimated medium samples. Fractions were tested for
activity by sandwich splat assay as described in example I.

The antifreeze activity in culture medium was purified by
gel permeation chromatography as follows.

- 10 The 14 cold acclimated 0.5M NaCl eluate from the Q
sepharose column (fraction 2) from above was acetone
precipitated and the pellet resuspended in 50µl 50mM
Tris/HCl + 0.15 M NaCl, pH 7.2. This was then centrifuged
at 10.000g for 10 minutes, and 20µl loaded onto a Superdex
15 75 gel permeation column on the Pharmacia SMART system. The
flow rate was 40µl/min and the mobile phase was 50mM
Tris/HCl + 0.15M NaCl, pH7.2. 80µl fractions were collected
and splatted. This procedure was repeated using t=14d non-
cold acclimated 0.5M NaCl eluate from the Q sepharose
20 column and fresh medium.

Further isolation of the active proteins can be done by SDS
PAGE analysis as described above.

Example VIII

Root extract from cold acclimatised carrot roots was prepared by scrubbing freshly pulled cold acclimatised carrots in cold water. The tops are removed and the juice extracted employing a domestic juice extractor (Russell Hobbs, model no 9915). The juice was frozen in 1 litre blocks and stored at -20°C prior to collection for use in ice cream trials.

10

The carrot AFP juice was added to the following ice cream formulation

INGREDIENT	parts by weight
Skimmed Milk Powder	10.000
Sucrose	13.000
MD40	4.000
Locust Bean Gum	0.144
Genulacta L100	0.016
MGP	0.300
Butteroil	8.000
Vanillin	0.012
Water	64.528
Carrot Extract (from cold acclimated carrots containing 1-10 mg AFP per kg)	4.472

25

Ice-cream was prepared by freezing above formulation and aeration to 106% overrun.

30

Measurements were made on fresh sample and on samples which had been abused by storage at -10 °C for a period of 10 days.

As a comparison a sample without carrot extract was measured in the same way. The measurements were done as follows:

- 5 Samples were equilibrated at -18 C in a Prolan Environmental cabinet for approximately 12 hours. Three samples were chosen representatively from each batch of ice cream and a slide was prepared from each in a Cryostat temperature control cabinet by smearing a thin layer of ice
10 cream from the centre of each block onto a microscopic slide. A single drop of white spirit was applied to the slide and a cover slip was then applied. Each slide, in turn, was then transferred to a temperature controlled microscope stage (Leit LaborLux S, Leica x10 objective,
15 temperature -18 °C). Images of ice-crystals (about 400 individual ice-crystals) were collected and relayed through a video camera (Sanyo CCD) to an image storage and analysis system (LEICA Q520MC).
- 20 The stored ice crystal images were highlighted manually by drawing around the perimeter which then highlights the whole crystal. Images of the highlighted crystals were then measured using the image analysis software which counts the number of pixels required to complete the longest straight
25 line (length), shortest straight line (breadth), the aspect ratio (length/breadth). The data for each individual ice crystal of a batch of ice cream was imported into a spreadsheet where analysis of the data set was carried out to find the mean, and standard deviation.
- 30
The ice Cream Hardness Measurements were carried out using a Hounsfield H10KM Universal Tester, a Hounsfield 100N Load Cell and a 10cm Cylindrical Stainless steel probe. The ice-cream samples were prepared by 16 Hour incubation of 486ml
35 ice cream blocks in a Prolan Temperature Control Cabinet set at -18 °C.

The ice cream block was removed from Prolan temperature control cabinet and placed the Hounsfield H10KM Universal Tester. The 10cm cylindrical probe was pushed into the ice cream block at a constant rate of 400mm/min to a depth of 20mm. The maximum force recorded during the compression was used and expressed as the ice cream Hardness. If cracking or brittle fracture of the sample was observed this was indicated in the right hand column

The following results were obtained

Sample	Ice Crystal Size Parameters				Material Properties	
	Mean Crystal Length / μm	Mean Crystal Breadth / μm	Mean Crystal Shape Factor / -	Mean Crystal Aspect Ratio / -	Hardness / N	Brittle Fracture observation
Carrot AFP - fresh	26.79 \pm 1.3	19.00 \pm 0.9	1.15 \pm 0.013	1.43 \pm 0.024	40.8	Yes
Carrot AFP - Abused	33.48 \pm 1.3	24.61 \pm 0.9	1.13 \pm 0.013	1.37 \pm 0.020	59.9	Yes
Cont.- Fresh	33.67 \pm 1.1	24.79 \pm 0.8	1.12 \pm 0.008	1.38 \pm 0.018	27.3	No
Cont.- Abused	61.77 \pm 2.7	46.54 \pm 2.0	1.11 \pm 0.010	1.37 \pm 0.020	32.7	No

The following conclusions can be drawn:

- a) Initial ice crystal size is smaller in ice cream containing Carrot AFP, thus carrot AFP is inhibiting recrystallization inhibition.
- b) Ice crystals in carrot AFP ice cream are retarded in their recrystallization processes.
- c) Ice crystal shape in carrot AFP ice creams are not

significantly different from crystal shapes seen in conventional ice creams.

- 5 d) Material properties of ice cream containing carrot AFP are modified from those noted for conventional ice cream. Namely, ice creams are harder than conventional ice cream but still softer than ice-cream containing e.g. fish AFPs. Secondly, ice cream containing carrot AFP was observed to fracture.

10

Similarly very good results can be obtained by using Geranium or Juncus squarrosus.

Example IX

This example describes the isolation of various proteins from winter rye and the testing thereof.

- 5 The leaves from 30 days cold acclimated rye plants were cut into 3 cm lengths and thororughly washed in distilled water to remove any cell contents. The leaf pieces were patted dry on paper towel and totally imersed in an extraction
- 10 medium of 5 mM EDTA, 10 mM ascorbic acid, 2 mM caproic acid, 2 mM benzamidine and 1 mM Phenylmethysulphonyl Fluoride (PMSF). They were then vacuum infiltrated in a Buchner flask for 60 minutes after which time the leaves were removed and patted completely dry. They were then
- 15 arranged lengthways is a cut off plastic syringe barrel and centrifuged gently at 2000 X g for 30 minutes. The apoplastic extract was collected in an eppendorf tube below the syringe.
- 20 The apoplastic extract was concentrated 7 times using and Amicon ultrafilter with a PM10 membrane. Initial purification was performed by loading 50 microliter of concentrated apoplastic extract onto a size exclusion. Superdex 75 PC 3.2/3.0 (separation range 3-70 kDa) column
- 25 on a SMART separation system, both from Pharmacia. The buffer was 50 mM Tris/HCl at pH 9.5. Separation was carried out at a flow rate of 50 microliter per minute and 50 microliter fractions were collected up to a volume of 2.5 ml and assayed for recrystallisation as described in
- 30 example I.

- Active fractions were loaded onto a strong anion exchange MonoQ FPLC column from Pharmacia, equilibrated in 50 mM Tris/HCl at pH 9.5, and the proteins were eluted using the
- 35 same buffer with an linear gradient to 0.5 M NaCl. The elution buffer was added to a concentration of 0.5 M NaCl over 25 minutes, held for 10 minutes and reduce to 0 M over

15 minutes. Chromatography was carried out at a flow rate of 1 ml per minute and 1 ml fractions were collected. The fractions which were positive in the test according to example I were concentrated on a Centricon PM10 centrifugal
5 concentrator at 7000 rps until the volume was reduced to 50 microlitre and loaded for a second time onto the S75 column. The fraction which satisfied the test of example i was a single peakd at approximately 150 mM salt.

- 10 This active fraction was separated on SDS PAGE (similar methodology as in example VII). This confirmed that the 32 kDa was the active fraction.

The 32 kDa fraction of winter rye protein can
15 advantageously be used in the preparation of frozen confectionery products.

Claims

1. A frozen confectionery products comprising one or more
5 AFPs derived from plants, wherein the AFPs in an
aqueous composition have an ice crystal size after
quick freezing to -40 °C or less, followed by storage
for 1 hour at -6 °C of less than 15 μ m.

2. A frozen confectionery product of claim 1, wherein the
10 AFP is derived from Polystichum mohriodes, Ranunculus
bitermatus, Nothofagus antartica, Cerastium fontanum,
Colobanthus quitensis, Rumex acetosella, Salix
fragilis, Calluna vulgaris, Aceana magellanica, Pisum
sativum, Acer saccharoides, Oxalis, Geranium, Daucus
15 carota (carrot), Vinca minor (periwinkle), Vinca
major, Polemonium, Buddleia, Forsythia, Sambucus
nigra, Juncus squarrosus, Carex aquatilis, Agrostis
tenuis, Deschampsia antartica, Festuca contracta,
Festuca rubra, Parodiochloa flabellata, Phleum
20 alpinum, Poa annua (speargrass), Poa pratensis
(Kentucky blue grass), Rostkovia magellanica,
Bambosoideae, Chorisodontium aciphyllum, Drepanocladus
uncinatus, Isothenicium myosuriodes, Polytrichum
alpestre, Alecatoria nigricans, Caloplaca regalis,
25 Himantormia lugubris, Hypogymnia physodes, Parmelia
subrudecta, Ramalina farinaceae, Stereocaulon glabrum,
Umbilicaria antartica, Usnea subfloridana, Poa
trivialis, Lolium perenne, Holcus lanatus, Bromus
sterilis and Festuca contracta.

30

3. Frozen confectionery product according to claim 2,
wherein the AFP is derived from the Lichen family, in
particular *Alectoria nigricans*, *Caloplaca regalis*,
Himantormia lugubris, *Hypogymnia physodes*, *Parmelia*
5 *subrudecta*, *Ramalina farinaceae*, *Stereocaulon glabrum*,
Umbilicaria antartica, *Usnea subfloridana*.
4. Frozen confectionery product according to claim 2,
wherein the AFP is derived from *Juncus squarrosus* or
10 *Geranium*.
5. Frozen confectionery product according to claim 2,
wherein the AFP maintain their ability to limit ice
crystal growth after heat treatment above a
15 temperature of 60°C for a period of at least 30
seconds, more preferred more than 1 minute.
6. Frozen confectionery product according to claim 5
wherein the AFP is derived from *Acer saccharoides*,
20 *Bamboo*, *Buddleia*, *Isothecium myosuroides*, *Ramalina*
farinaceae, *Usnea subfloridana*, *Forsythia*, *Oxalis*, *Poa*
trivialis, *Lolium perenne*, *Holcus lanatus*, *Bromus*
sterilis, *Parodiochloa flabellata*, *Deschampsia*
25 *antartica*, *Carex aquatilis*, *Colobanthus quintensis* and
Agrostis tenuis, *Festuca contracta*, *Poa annua*.
7. Frozen confectionery product according to claim 1,
wherein the AFP is derived from a non-toxic plant.
- 30 8. Frozen confectionery product according to claim 2,
wherein the AFP is the 32 kDa protein derived from
winter rye.
9. Frozen confectionery product according to claim 2,
35 wher in the product is substantially free from AFPs
from winter rye.



Application No: GB 9714412.5
Claims searched: 1 to 9

Examiner: S J Pilling
Date of search: 30 September 1997

Patents Act 1977
Search Report under Section 17

Databases searched:

UK Patent Office collections, including GB, EP, WO & US patent specifications, in:
UK Cl (Ed.O): C3H (HF1, HFX)
Int Cl (Ed.6): C07K 14/415, A23G 9/00, 9/02, A23L 3/36
Other: ONLINE: WPI, BIOTECH

Documents considered to be relevant:

Category	Identity of document and relevant passage	Relevant to claims
X	WO 92/22581 A1 (UNIVERSITY OF WATERLOO) see page 8 lines 3 to 12, page 14 lines 23 to 28.	1,7
X	US 4565643 (ARAI & WATANABE) see column 3 lines 36 to 40, column 3 line 64 to column 4 line 2 and Comparative Example 2.	1,7
X	Biotechnology Advances, Vol. 13, No. 3, 1995, M Griffith <i>et al</i> "Antifreeze proteins and their potential use in frozen foods", pages 375 to 402	1,7,9
A	Plant Physiol., Vol. 104, 1994, W C Hon <i>et al</i> "Extraction and isolation of antifreeze proteins from Winter Rye (<i>Secale cereale</i>) leaves" pages 971 to 980.	-
A	Cryobiology, Vol. 30, 1993, J G Duman & T M Olsen, "Thermal hysteresis protein activity in bacteria, fungi and phylogenetically diverse plants", pages 322 to 328.	-
A	Advances in Low-Temperature Biology, Vol. 2, 1993, J G Duman <i>et al</i> , "Thermal hysteresis proteins", pages 131 to 182.	-

X	Document indicating lack of novelty or inventive step	A	Document indicating technological background and/or state of the art.
Y	Document indicating lack of inventive step if combined with one or more other documents of same category.	P	Document published on or after the declared priority date but before the filing date of this invention.
&	Member of the same patent family	E	Patent document published on or after, but with priority date earlier than, the filing date of this application.



The Patent Office

32

Application No: GB 9714411.7
Claims searched: 1 to 12

Examiner: S J Pilling
Date of search: 26 September 1997

Category	Identity of document and relevant passage	Relevant to claims
X	Biochemical and Biophysical Research Communications, Vol. 183, No. 3, March 1992, C Lin & M F Thomashow, " <i>A cold-regulated Arabidopsis gene encodes a polypeptide having potent cryoprotective activity</i> ", pages 1103 to 1108.	Claims 1 and 3 at least
A	Protein Engineering, Vol. 4, No. 8, 1991, X Li <i>et al</i> " <i>Expression and characterization of an active and thermally more stable recombinant antifreeze polypeptide from ocean pout, Macrozoarces americanus, in Escherichia coli: improved expression by the modification of the secondary structure of the mRNA</i> ", pages 995 to 1002	
X	Biochimica et Biophysica Acta, Vol. 412, 1975, H G Volger & U Hueber " <i>Cryoprotective leaf proteins</i> ", pages 335 to 349	Claims 1 and 3 at least

X	Document indicating lack of novelty or inventive step	A	Document indicating technological background and/or state of the art.
Y	Document indicating lack of inventive step if combined with one or more other documents of same category.	P	Document published on or after the declared priority date but before the filing date of this invention.
&	Member of the same patent family	E	Patent document published on or after, but with priority date earlier than, the filing date of this application.

